

A Bounded Divergence Measure Based on The Bhattacharyya Coefficient

Shivakumar Jolad, Ahmed Roman, and Mahesh C. Shastry, *Member, IEEE*,

Abstract—We introduce a new divergence measure, the bounded Bhattacharyya distance (BBD), for quantifying the dissimilarity between probability distributions. BBD is based on the Bhattacharyya coefficient (fidelity), and is symmetric, positive semi-definite, and bounded. Unlike the Kullback-Leibler divergence, BBD does not require probability density functions to be absolutely continuous with respect to each other. We show that BBD belongs to the class of Csiszar f-divergence and derive certain relationships between BBD and well known measures such as Bhattacharyya, Hellinger and Jensen-Shannon divergence. Bounds on the Bayesian error probability are established with BBD measure. We show that the curvature of BBD in the parameter space of families of distributions is proportional to the Fisher information. For distributions with vector valued parameters, the curvature matrix can be used to obtain the Rao geodesic distance between them. We also discuss a potential application of probability distance measures in model selection.

Index Terms—Signal detection, Bhattacharyya distance, divergence, dissimilarity measure, f-divergence, error probability.

I. INTRODUCTION

Divergence measures for the distance between two probability distributions have been extensively studied in the last six decades [1], [2], [3], [4], [5]. These measures are widely used in varied fields such as pattern recognition [6], [7], [8], signal detection [9], [10], Bayesian model validation [11] and quantum information theory [12], [13]. Distance measures try to achieve two main objectives (which are not mutually exclusive): to assess (1) how “close” two distributions are compared to others and (2) how “easy” it is to distinguish between one pair than the other [2].

There is a plethora of distance measures available to assess the convergence (or divergence) of probability distributions. Many of these measures are not metrics in the strict sense, as they may not satisfy either the symmetry of arguments or triangle inequality. In applications, the choice of the measure depends on the interpretation of the metric in terms of the problem considered, its analytical properties and ease of computation [14]. One of the most well-known divergence measures is the Kullback-Leibler divergence (KLD)[1], [4]. Although it is widely used, KLD can create problems in specific applications. Specifically, it is unbounded above and requires that the distributions be ‘absolutely continuous’ with respect to each other. Various other information theoretic

measures have been introduced keeping in view ease of computation and utility in problems of signal selection and pattern recognition. Of these measures, Bhattacharyya distance [15], [9], [16] and Chernoff distance [17], [6], [18] have been widely used in signal processing. However, these measures are again unbounded from above. Many bounded divergence measures such as Variational, Hellinger distance [6], [19] and Jensen-Shannon metric [20], [21], [22] have been studied extensively. However, utility of these measures varies depending on properties such as tightness of bounds on error probabilities, information theoretic interpretation, and generalization to multiple probability distributions. Here we introduce a bounded measure based on the Bhattacharyya coefficient which shares a close relationship with Hellinger and Jensen-Shannon divergences. Our bounded Bhattacharyya distance (BBD) measure belongs to the class of f-divergences and thus inherits all its general properties. We prove an extension of Bradt-Karlin theorem for BBD which shows the existence of prior probabilities for which ranking of divergence is mapped to the ranking of Bayes error probabilities. Based on Bhattacharyya coefficient we also show upper and lower bounds on the error probabilities. We show that BBD can be expressed as f-divergence measure. For many applications in Biology, more than two probability measures have to be distinguished from each other. Following Rao [21] and Lin [22] we introduce a generalized BBD measure for a generic set of probability distributions. We also propose a potential application of distance measure to model selection in this work.

Our paper is organized as follows: Section I is the current introduction. In Section II, we discuss the well known divergence measures Kullback-Leibler and Bhattacharyya and introduce our bounded measure. In Section III, we derive several interesting properties of our measure such as positive semi-definiteness, relation with Hellinger and Jensen-Shannon metric and utility for computing probability of error. Generalized BBD measures are discussed in Section IV. In Section V, the relation with Fisher information and Rao differential metric is derived. In the final section, we propose a method for using probability distance measures for goodness of fit and model selection. In the Appendix we provide the expressions for BBD measures for some commonly used distributions.

II. DIVERGENCE MEASURES

In the following subsection we consider a measurable space Ω with σ algebra $\mathcal{B}$ and the set of all probability measures $\mathcal{M}$ on $(\Omega, \mathcal{B})$. Let P and Q denote probability measures on $(\Omega, \mathcal{B})$

Shivakumar Jolad is currently with the Indian Institute of Technology - Gandhinagar, Ahmedabad, India e-mail: shiva.jolad@iitgn.ac.in. Major part of the work was done when S.J. was working with Ahmed Roman at the Department of Physics, Virginia Tech, Blacksburg, VA 24061, USA e-mail: mido@vt.edu

Mahesh Shastry is with the Department of Electrical Engineering, The Pennsylvania State University, PA 16803 USA e-mail: mcs312@psu.edu

and p and q denote their densities with respect to a common measure μ . We recall the definition of absolute continuity [Cite textbook source, and use consistent notations]:

Absolute Continuity A measure P on the Borel subsets of the real line is absolutely continuous with respect to Lebesgue measure Q , if $P(A) = 0$, for every Borel subset $A \in \mathcal{B}$ for which $Q(A) = 0$, and is denoted by $P \ll Q$.

A. Kullback-Leibler divergence

The Kullback-Leibler divergence (KLD) (or relative entropy) [1], [4] between two distributions P, Q with densities $p(x)$ and $q(x)$ is given by:

$$I(P, Q) \equiv \int p(x) \log \left(\frac{p(x)}{q(x)} \right) dx. \quad (1)$$

The symmetrized version is given by $I_{\text{symm}}(P, Q) \equiv (I(P, Q) + I(Q, P))/2$ [9], $I(P, Q) \in [0, \infty]$. It diverges if $\exists x_0 : q(x_0) = 0$ and $p(x_0) \neq 0$.

KLD is defined only when P is absolutely continuous w.r.t. Q . This feature can be problematic in numerical computations when the measured distribution has zero values.

B. Bhattacharyya Distance

Bhattacharyya distance is a widely used measure in signal selection and pattern recognition [9]. It is defined as:

$$B(P, Q) \equiv -\ln \left(\int \sqrt{p(x)q(x)} dx \right) = -\ln(\rho), \quad (2)$$

where the term in parenthesis $\rho(P, Q) \equiv \int \sqrt{p(x)q(x)} dx$ is called Bhattacharyya coefficient [23], [15] in pattern recognition, affinity in theoretical statistics, and fidelity in quantum information theory. Unlike in the case of KLD, the Bhattacharyya distance avoids the requirement of absolute continuity. Its a special case of Chernoff distance

$$C_\alpha(P, Q) \equiv -\ln \left(\int p^\alpha(x) q^{1-\alpha}(x) dx \right),$$

with $\alpha = 1/2$. For discrete probability distributions, $\rho \in [0, 1]$ is interpreted as a scalar product of the probability vectors $\mathbf{P} = (\sqrt{p_1}, \sqrt{p_2}, \dots, \sqrt{p_n})$ and $\mathbf{Q} = (\sqrt{q_1}, \sqrt{q_2}, \dots, \sqrt{q_n})$. Bhattacharyya distance is symmetric, positive-semidefinite, and unbounded ($0 \leq B \leq \infty$). It is finite as long as there exists some region $S \subset X$ such that whenever $x \in S : p(x)q(x) \neq 0$.

C. Bounded Bhattacharyya distance measure

In many applications, in addition to the desirable properties of the Bhattacharyya distance, boundedness is required. We propose a new bounded measure of Bhattacharyya distance as below,

$$\zeta(P, Q) \equiv -\log_2 \left[\frac{1 + \int \sqrt{p(x)q(x)} dx}{2} \right] = -\log_2 \left[\frac{1 + \rho}{2} \right]. \quad (3)$$

With the choice of base 2 for the logarithm, we normalize the maximum value to 1. For convenience, we shall also refer to




Fig. 1. Comparison of Hellinger and bounded Bhattacharyya distance measures ζ and ξ .

the bounded Bhattacharyya measure as the *zeta* measure, and use the terms BBD and ζ interchangeably. The term under the logarithm in Eq. 3 is the average of complete and actual overlaps of two distributions. From this measure, we can form another closely related measure by taking the square root of the ζ :

$$\xi(P, Q) \equiv \sqrt{\zeta(P, Q)}. \quad (4)$$

This measure is also symmetric, positive definite and bounded between $[0, 1]$. We will not delve into the properties of ξ measure in this paper.

Another widely used bounded measure based on the Bhattacharyya distance is the Hellinger distance [6], [9], [24], [25]

$$H(P, Q) \equiv \sqrt{1 - \rho(P, Q)}. \quad (5)$$

We note that $H \in [0, 1]$ and is concave in ρ ($\partial_\rho^2 H = -1/2(1-\rho)^{3/2} < 0$). In contrast our ζ measure is convex in ρ ($\partial_\rho^2 \zeta = 1/(1+\rho)^2 \log 2 > 0$), where as ξ is neither concave nor convex. A comparison between Hellinger and our ζ and ξ measures as function of Bhattacharyya coefficient is given in Fig. 1.

III. PROPERTIES OF ζ AND ξ MEASURES

Theorem III.1 (Positive semi-definite). *ζ measure is symmetric, positive semi-definite and bounded in the interval $[0, 1]$.*

Proof: By using arithmetic and geometric means inequality, we obtain the following:

$$\begin{aligned} \int \sqrt{p(x)q(x)} dx &\leq \int \left(\frac{p(x) + q(x)}{2} \right) dx \\ &= \frac{1}{2} \int (p(x) + q(x)) dx = 1. \end{aligned} \quad (6)$$

Which leads to :

$$0 \leq -\log_2 \left[\frac{1 + \int \sqrt{p(x)q(x)} dx}{2} \right] \leq 1. \quad (7)$$

and hence

$$0 \leq \zeta(P, Q) \leq 1. \quad (8)$$

Special cases:

$$\zeta(P, Q) = \begin{cases} 0 & \text{if } P = Q \text{ almost everywhere} \\ 1 & \text{if } p(x)q(x) = 0 \text{ for } \forall x \in X \\ (0, 1) & \text{otherwise} \end{cases} \quad (9)$$

ζ is symmetric by inspection:

$$\zeta(P, Q) = \zeta(Q, P). \quad (10)$$

Error probability: The optimal Bayes error probabilities (see eg: [7], [26]) for classifying two events P_1, P_2 with densities $p_1(x)$ and $p_2(x)$ with prior probabilities $\Gamma = \{\pi_1, \pi_2\}$ is given by

$$P_e = \int \min[\pi_1 p_1(x), \pi_2 p_2(x)] dx. \quad (11)$$

Let $p_i(x)$ ($i = 1, 2$) be parameterized by α or β . In signal detection literature, a signal set α is considered better than set β for the densities $p_i(x)$ ($i = 1, 2$), when the error probability is less for α than for β [9].

We can also rank the parameters by means of some divergence D . That is, we can say the set α is better (in the divergence say) than set β , if $D_\alpha(P_1, P_2)$ is larger than $D_\beta(P_1, P_2)$. In general, it is not true that if $D_\alpha(P_1, P_2) > D_\beta(P_1, P_2)$, then $P_e(\alpha) < P_e(\beta)$. Bradt and Karlin proved the following theorem relating error probabilities and symmetric KLD:

Theorem III.2 (Bradt and Karlin [27]). *If $J_\alpha(P_1, P_2) > J_\beta(P_1, P_2)$, then $\exists$ a set of prior probabilities $\Gamma = \{\pi_1, \pi_2\}$ for two hypothesis g_1, g_2 , for which*

$$P_e(\alpha, \Gamma) < P_e(\beta, \Gamma) \quad (12)$$

where $P_e(\alpha, \Gamma)$ is the error probability with parameter α and prior probability Γ .

It is clear that the theorem asserts existence, but no method of finding these prior probabilities. Kailath [9] proved the applicability of Bradt-Karlin Theorem for Bhattacharyya distance measure. We follow the same route and show that the ζ -measure satisfies a similar property using the following theorem by Blackwell.

Theorem III.3 (Blackwell [28]). *$P_e(\beta, \Gamma) \leq P_e(\alpha, \Gamma)$ for all prior probabilities Γ if and only if*

$$\mathbb{E}_\beta[\Phi(L_\beta)|g^{(2)}] \leq \mathbb{E}_\alpha[\Phi(L_\alpha)|g^{(2)}],$$

$\forall$ continuous concave functions $\Phi(L)$, where $L_\omega = p_1(x, \omega)/p_2(x, \omega)$ is the likelihood ratio with $\omega = \{\alpha, \beta\}$ and $\mathbb{E}_\omega[\Phi(L_\omega)|g^{(2)}]$ is the expectation of $\Phi(L_\omega)$ under the hypothesis $g^{(2)}$.

Theorem III.4. *If $\zeta(\alpha) > \zeta(\beta)$, or equivalently $\rho(\alpha) < \rho(\beta)$ then $\exists$ a set of prior probabilities $\Gamma = \{\pi_1, \pi_2\}$ for two hypothesis g_1, g_2 , for which*

$$P_e(\alpha, \Gamma) < P_e(\beta, \Gamma). \quad (13)$$

Proof: The proof closely follows Kailath [9]. First note that $\sqrt{L}$ is a concave function of L (likelihood ratio), and

$$\begin{aligned} \rho(\alpha) &= \sum_{x \in X} \sqrt{p_1(x, \alpha)p_2(x, \alpha)} \\ &= \sum_{x \in X} \sqrt{\frac{p_1(x, \alpha)}{p_2(x, \alpha)}} p_2(x, \alpha) \\ &= \mathbb{E}_\alpha[\sqrt{L_\alpha}|g^{(2)}]. \end{aligned} \quad (14)$$

Similarly

$$\rho(\beta) = \mathbb{E}_\beta[\sqrt{L_\beta}|g^{(2)}] \quad (15)$$

Hence, $\rho(\alpha) < \rho(\beta) \Rightarrow$

$$\mathbb{E}_\alpha[\sqrt{L_\alpha}|g^{(2)}] < \mathbb{E}_\beta[\sqrt{L_\beta}|g^{(2)}]. \quad (16)$$

Suppose assertion of the stated theorem is not true, then for all Γ , $P_e(\beta, \Gamma) \leq P_e(\alpha, \Gamma)$. Then by Theorem III.3, $\mathbb{E}_\beta[\Phi(L_\beta)|g^{(2)}] \leq \mathbb{E}_\alpha[\Phi(L_\alpha)|g^{(2)}]$ which contradicts our result in Eq. 16. ■

Theorem III.5. $0 \leq \zeta(P, Q) \leq \xi(P, Q) \leq H(P, Q)$.

Proof: We have already shown that $0 \leq \zeta \leq 1$ in Theorem III.1. Since $\xi = \sqrt{\zeta}$, it follows that $0 \leq \zeta(P, Q) \leq \xi(P, Q)$. We use the generalized Bernoulli inequality

$$(1+x)^r \leq 1+rx, \quad 0 \leq r \leq 1, \quad x > -1. \quad (17)$$

Set $x = 1$ and $r = \rho = \int \sqrt{p(x)q(x)} dx$, the Bhattacharyya coefficient. Hence

$$\begin{aligned} 0 &\leq (1+1)^\rho = 2^\rho \leq 1+\rho \leq 2 \\ 0 &\leq \rho \leq \log_2(1+\rho) \leq 1 \\ 1 &\geq \sqrt{1-\rho} \geq \sqrt{1-\log_2(1+\rho)} \geq 0 \\ 1 &\geq \sqrt{1-\rho} \geq \sqrt{-\log_2 \frac{(1+\rho)}{2}} \geq 0. \end{aligned} \quad (18)$$

Hence we get

$$1 \geq H \geq \xi \geq 0. \quad (19)$$

Theorem III.6.

$$\xi \leq H \leq \sqrt{\ln 4} \xi \quad (20)$$

where 1 and $\sqrt{\ln 4}$ are sharp.

Proof: Sharp lower bound has been proved in Theorem III.5. Sharpest upper bound is achieved via taking $\sup_{\rho \in [0,1]} \frac{H(\rho)}{\xi(\rho)}$. Define

$$f(\rho) \equiv \frac{H(\rho)}{\xi(\rho)} = \frac{\sqrt{1-\rho}}{\sqrt{-\log_2(1+\rho)/2}}, \quad (21)$$

$$g(\rho) \equiv f^2(\rho). \quad (22)$$

We note that $g(\rho)$ is continuous and has no singularities whenever $\rho \in [0, 1)$. Hence

$$g'(\rho) = \frac{\frac{1-\rho}{1+\rho} + \ln(\frac{1+\rho}{2})}{\ln^2 \frac{\rho+1}{2}} \ln 2 \geq 0.$$

It follows that if $f(\rho)$ is non-decreasing and hence $\sup_{\rho \in [0,1]} g(\rho) = \lim_{\rho \rightarrow 1} g(\rho) = \ln(4)$. Thus

$$f(\rho) \leq \sqrt{\sup_{\rho \in [0,1]} g(\rho)} = \sqrt{\ln(4)}. \quad (23)$$

Hence

$$H/\xi \leq \sqrt{\ln 4}. \quad (24)$$

■

Jensen-Shannon Divergence: The Jensen difference between two distributions P_1, P_2 , with densities $p_1(x), p_2(x)$ and weights (λ_1, λ_2) ; $\lambda_1 + \lambda_2 = 1$, is defined as,

$$\mathcal{J}_{\lambda_1, \lambda_2}(P_1, P_2) = H(\lambda_1 p_1 + \lambda_2 p_2) - \lambda_1 H(p_1) - \lambda_2 H(p_2). \quad (25)$$

Jensen-Shannon divergence (JSD) [20], [21], [22] is based on the Jensen difference and is given by:

$$\begin{aligned} JS(P, Q) &= \mathcal{J}_{1/2, 1/2}(P, Q) \\ &= \frac{1}{2} \int \left[p(x) \log \left(\frac{2p(x)}{p(x) + q(x)} \right) \right. \\ &\quad \left. + q(x) \log \left(\frac{2q(x)}{p(x) + q(x)} \right) \right] dx. \end{aligned} \quad (26)$$

The structure and goals of JSD and BBD measures are similar. The following theorem compares the two metrics using Jensen's inequality.

Lemma III.7. Jensen's Inequality: For a convex function ψ , $\mathbb{E}[\psi(X)] \geq \psi(\mathbb{E}[X])$.

Theorem III.8 (Relation to Jensen-Shannon measure). $JS(P, Q) \geq \frac{2}{\log 2} \zeta(P, Q) - \log 2$

We use the un-symmetrized Jensen-Shannon metric for the proof.

Proof:

$$\begin{aligned} JS(P, Q) &= \int p(x) \log \frac{2p(x)}{p(x) + q(x)} dx \\ &= -2 \int p(x) \log \frac{\sqrt{p(x)} + \sqrt{q(x)}}{\sqrt{2p(x)}} dx \\ &\geq -2 \int p(x) \log \frac{\sqrt{p(x)} + \sqrt{q(x)}}{\sqrt{2p(x)}} dx \\ &\quad (\text{since } \sqrt{p(x)} + \sqrt{q(x)} \leq \sqrt{p(x)} + \sqrt{q(x)}) \\ &= \mathbb{E}_P \left[-2 \log \frac{\sqrt{p(X)} + \sqrt{q(X)}}{\sqrt{2p(X)}} \right] \\ &\geq -2 \log \mathbb{E}_P \left[\frac{\sqrt{p(X)} + \sqrt{q(X)}}{\sqrt{2p(X)}} \right] \\ &\quad (\text{by Jensen's inequality}) \\ &\quad \mathbb{E}[-\log f(X)] \geq -\log \mathbb{E}[f(X)] \\ &= -2 \log \int p(x) \frac{(\sqrt{p(x)} + \sqrt{q(x)})}{\sqrt{2p(x)}} dx \\ &= -2 \log \int \sqrt{p(x)} \left(\frac{\sqrt{p(x)} + \sqrt{q(x)}}{2} \right) \sqrt{2} dx \end{aligned}$$

$$\begin{aligned} &= -2 \log \left(\frac{1 + \int \sqrt{p(x)q(x)}}{2} \right) - \log 2 \\ &= 2 \left(\frac{\zeta(p(x), q(x))}{\log 2} \right) - \log 2 \\ &= \frac{2}{\log 2} \zeta(P, Q) - \log 2. \end{aligned} \quad (27)$$

Therefore we have the result,

$$JS(P, Q) \geq \frac{2}{\log 2} \zeta(P, Q) - \log 2. \quad (28)$$

■

A. Bounds on Error Probability

Error probabilities are hard to calculate in general. Tight bounds on P_e are often extremely useful in practice. Kailath [9] has shown bounds on P_e in terms of the Bhattacharyya coefficient ρ :

$$\frac{1}{2} \left[2\pi_1 - \sqrt{1 - 4\pi_1\pi_2\rho^2} \right] \leq P_e \leq \left(\pi_1 - \frac{1}{2} \right) + \sqrt{\pi_1\pi_2}\rho, \quad (29)$$

with $\pi_1 + \pi_2 = 1$. If the priors are equal $\pi_1 = \pi_2 = 1/2$, the expression simplifies to

$$\frac{1}{2} \left[1 - \sqrt{1 - \rho^2} \right] \leq P_e \leq \frac{1}{2} \rho. \quad (30)$$

Substituting $\rho = 2^{1-\zeta} - 1$, we can get the bounds in terms of our ζ measure. For the equal prior probabilities case, Bhattacharyya coefficient gives a tight upper bound for large systems when $\rho \rightarrow 0$ (zero overlap) and the observations are independent and identically distributed. These bounds are also useful to discriminate between two processes with arbitrarily low error probability [9].

B. f-divergence

A class of divergence measures called f-divergences were introduced by Csiszar [29], [30] and independently by Ali and Silvey [2] (see [6] for review). It encompasses many well known divergence measures including KLD, variational and Bhattacharyya distance. In this section, we show that our ζ measure belongs to the class of f-divergences.

f-divergence [6] Consider a measurable space Ω with σ algebra $\mathcal{B}$. Let λ be a measure on $(\Omega, \mathcal{B})$ such that any probability laws P_1 and P_2 are absolutely continuous with respect to λ , with densities p_1 , and p_2 . Let $\tilde{f}$ be a continuous *convex* real function on $\mathbb{R}^+$, and let g be an *increasing* function on $\mathbb{R}$. The class of divergence coefficients between two probabilities:

$$d(P_1, P_2) = g \left[\mathbb{E}_1 \left\{ \tilde{f} \left(\frac{p_2}{p_1} \right) \right\} \right] \quad (31)$$

are called the f-divergence measure w.r.t. functions $(\tilde{f}, g)$. Here $p_2/p_1 = L$ is the likelihood ratio and $\mathbb{E}_1$ is the expectation w.r.t. to P_1 .

The $\zeta(P_1, P_2)$ metric can be written as the following f divergence:

$$\tilde{f}(x) = -\frac{1}{4}(1 + \sqrt{x})^2, \quad g(F) = -\log_2(-F), \quad (32)$$

where,

$$\begin{aligned}
F &= \mathbb{E}_1 \left[\tilde{f} \left(\frac{p_2}{p_1} \right) \right] \\
&= - \int p_1 \frac{1}{4} \left(1 + \sqrt{\frac{p_2}{p_1}} \right)^2 d\lambda \\
&= - \frac{1}{4} \int (\sqrt{p_1} + \sqrt{p_2})^2 d\lambda \\
&= - \frac{1 + \rho}{2}.
\end{aligned} \tag{33}$$

and

$$g(F) = -\log_2(-F) = -\log_2 \left(\frac{1 + \rho}{2} \right) = \zeta(P_1, P_2). \tag{34}$$

C. Generalized ζ measure

In decision problems involving more than two classes it is very useful to have divergence measures involving more than two distributions [31], [22]. We use the generalized geometric mean (GGM) concept to define ζ divergence for more than two distributions. The GGM of n variables $p_1, p_2, \dots, p_n$ with weights $\alpha_1, \alpha_2, \dots, \alpha_n$, such that $\alpha_i \geq 0$, $\sum_i \alpha_i = 1$, is given by

$$GGM(\{p_i\}) = \prod_{i=1}^n p_i^{\alpha_i}.$$

The generalized version of ζ measure for n probability measures $P_1, P_2, \dots, P_n$ can be defined as:

$$\zeta_{\alpha_1, \dots, \alpha_n} = -\log \left(\frac{1 + \int_X \prod_{i=1}^n p_i^{\alpha_i} d\lambda}{2} \right), \tag{35}$$

where $\alpha_i \geq 0$, $\sum_{i=1}^n \alpha_i = 1$. We note that,

$$0 \leq \zeta_{\alpha_1, \dots, \alpha_n} \leq 1,$$

since the weighted geometric mean is maximized when all the p_i 's are the same, and is minimized when any two of the probability densities p_i 's are perpendicular to each other.

IV. CURVATURE OF ζ METRIC

In statistics, the information that an observable random variable X carries about an unknown parameter θ (on which it depends) is given by the Fisher information. One of the important properties of f-divergence of two distributions of the same parametric family is that their curvature measures the Fisher information. Following the approach pioneered by Rao [32], in this section, we relate the curvature of ζ measures to the Fisher information and derive the differential curvature metric. The following discussions closely follow DasGupta [19].

Definition Let $\{f(x|\theta); \theta \in \Theta \subseteq \mathbb{R}\}$, be a family of densities indexed by real parameter θ , with some regularity conditions ($f(x|\theta)$ is absolutely continuous).

$$\zeta(\theta, \phi) = -\log_2 \left(\frac{1 + \rho(\theta, \phi)}{2} \right) = Z_\theta(\phi) \tag{36}$$

where $\rho(\theta, \phi) = \int \sqrt{f(x|\theta)f(x|\phi)} dx$

Theorem IV.1. Curvature of $Z_\theta(\phi)|_{\phi=\theta}$ is the Fisher information of $f(x|\theta)$ up to a multiplicative constant.

Proof: Expand $Z_\theta(\phi)$ around theta

$$\begin{aligned}
Z_\theta(\phi) &= Z_\theta(\theta) + (\phi - \theta) \frac{d}{d\phi} Z_\theta(\phi) \Big|_{\phi=\theta} \\
&\quad + \frac{(\phi - \theta)^2}{2} \frac{d^2}{d\phi^2} Z_\theta(\phi) \Big|_{\phi=\theta} + \dots
\end{aligned} \tag{37}$$

Let us observe some properties of Bhattacharyya coefficient $\rho(\theta, \phi)$

$$\begin{aligned}
\rho(\theta, \phi) &= \rho(\phi, \theta) \\
\rho(\theta, \theta) &= 1.
\end{aligned} \tag{38}$$

and its derivatives:

$$\begin{aligned}
\frac{\partial \rho(\theta, \phi)}{\partial \phi} \Big|_{\phi=\theta} &= \frac{1}{2} \frac{\partial}{\partial \theta} \int f(x|\theta) dx = 0, \\
\frac{\partial^2 \rho(\theta, \phi)}{\partial \phi^2} \Big|_{\phi=\theta} &= \frac{-1}{4} \int \frac{1}{f(x|\theta)} \left(\frac{\partial f}{\partial \theta} \right)^2 dx \\
&\quad + \frac{1}{2} \frac{\partial^2}{\partial \theta^2} \int f(x|\theta) dx \\
&= \frac{-1}{4} \int f(x|\theta) \left(\frac{\partial \log f(x|\theta)}{\partial \theta} \right)^2 dx + 0 \\
&= \frac{-1}{4} I_f(\theta).
\end{aligned} \tag{39}$$

where $I_f(\theta)$ is the Fisher Information of distribution $f(x|\theta)$

$$I_f(\theta) = \int f(x|\theta) \left(\frac{\partial \log f(x|\theta)}{\partial \theta} \right)^2 dx. \tag{40}$$

Using the above relationships, we can write down the terms in the expansion of Eq. 37 (neglecting $\log 2$ factor for brevity)

$$\begin{aligned}
Z_\theta(\theta) &= 1 \\
\frac{\partial Z_\theta(\phi)}{\partial \phi} \Big|_{\phi=\theta} &= \frac{-1}{1 + \rho} \frac{\partial \rho}{\partial \phi} \Big|_{\phi=\theta} = 0 \\
\frac{\partial^2 Z_\theta(\phi)}{\partial \phi^2} &= \frac{1}{(1 + \rho)^2} \left(\frac{\partial \rho}{\partial \phi} \right)^2 - \frac{1}{1 + \rho} \frac{\partial^2 \rho}{\partial \phi^2} \\
\frac{\partial^2 Z_\theta(\phi)}{\partial \phi^2} \Big|_{\phi=\theta} &= 0 - \frac{1}{1 + \rho(\theta, \theta)} \frac{-I_f(\theta)}{4} \\
&= \frac{1}{8} I_f(\theta),
\end{aligned} \tag{41}$$

We can relate the ζ metric to the Fisher Information as (converting back to base 2 units)

$$Z_\theta(\phi) = \frac{(\phi - \theta)^2}{2} \frac{1}{8 \log 2} I_f(\theta) + \dots \tag{42}$$

■

A. Differential Metrics

Rao [33] generalized the Fisher information to multivariate densities with vector valued parameters to obtain a “geodesic” distance between two parametric distributions P_θ, P_ϕ of the same family (see 15.4.2 in A DasGupta [19] for details). We derive such a metric for ζ measure.

Let $\theta, \phi \in \Theta \subseteq \mathbb{R}^p$, then using the fact that

$$\left. \frac{\partial Z(\theta, \phi)}{\partial \theta_i} \right|_{\phi=\theta} = 0, \quad (43)$$

we can easily show that

$$\begin{aligned} dZ_\theta &= \sum_{i,j=1}^p \frac{\partial^2 Z_\theta}{\partial \theta_i \partial \theta_j} d\theta_i d\theta_j + \dots, \\ &= \sum_{i,j=1}^p g_{ij} d\theta_i d\theta_j + \dots \end{aligned} \quad (44)$$

The curvature metric g_{ij} can be used to find the geodesic on the curve $\eta(t)$, $t \in [0, 1]$ with

$$\mathcal{C} = \eta(t) : \quad \eta(0) = \theta \quad \eta(1) = \phi, \quad (45)$$

The geodesic distance between θ and ϕ can be obtained by minimizing the length

$$s = \int_0^1 \sqrt{g_{ij}(\eta) \dot{\eta}_i \dot{\eta}_j} dt, \quad (46)$$

with the constraints in Eq. 45. The geodesic equation to be solved (assuming summation convention) are:

$$\frac{d^2 \eta_k}{dt^2} + \Gamma_{ijk} \frac{d\eta_i}{dt} \frac{d\eta_j}{dt} = 0, \quad (47)$$

where the Christoffel tensor Γ_{ijk} is given by:

$$\Gamma_{ijk} = \frac{1}{2} \left[\frac{\partial g_{jk}}{\partial \theta_i} + \frac{\partial g_{ki}}{\partial \theta_j} - \frac{\partial g_{ij}}{\partial \theta_k} \right].$$

For our $\zeta(\theta, \phi)$ metric, one can easily show (following steps in Theorem IV.1) the following result (neglecting $\log 2$ factor for brevity):

$$\begin{aligned} \left. \frac{\partial \zeta}{\partial \phi_i} \right|_{\phi=\theta} &= -\frac{1}{1+\rho} \left. \frac{\partial \rho}{\partial \phi_i} \right|_{\phi=\theta} = 0. \\ \left. \frac{\partial^2 \zeta}{\partial \phi_i \partial \phi_j} \right|_{\phi=\theta} &= \frac{1}{(1+\rho)^2} \left. \frac{\partial \rho}{\partial \phi_i} \frac{\partial \rho}{\partial \phi_j} \right|_{\phi=\theta} - \frac{1}{1+\rho} \left. \frac{\partial^2 \rho}{\partial \phi_i \partial \phi_j} \right|_{\phi=\theta} \\ &= \frac{1}{8} \mathbb{E}_f \left(\frac{\partial f(x|\theta)}{\partial \theta_i} \frac{\partial f(x|\theta)}{\partial \theta_j} \right) \end{aligned} \quad (48)$$

This is just a numerical factor times the Fisher information metric for KLD. This is due to the fact that curvature metric of all Csiszar f-divergences are just scalar multiple KLD measure [19], [6]:

$$g_{ij}^{\tilde{f}}(\theta) = \tilde{f}''(1) g_{ij}(\theta). \quad (49)$$

For our ζ metric

$$\begin{aligned} \tilde{f}''(x) &= \left(-\frac{1}{4}(1+\sqrt{x})^2 \right)'' \\ &= \frac{1}{8x^{3/2}} \\ \tilde{f}''(1) &= 1/8. \end{aligned} \quad (50)$$

Hence the result. It follows that the geodesic distance for our metric is same KLD geodesic distance up to a multiplicative factor. KLD geodesic distances are tabulated in DasGupta [19].

V. MODEL SELECTION

Model selection and parameter estimation of models of observed data are common problems in statistics, social science, life science and engineering. Many techniques exists for parameter estimation such as mean square error, maximum likelihood estimators, and Bayes estimators. We propose a method which can be potentially used to estimate parameters by minimizing the distance between the observed distribution and model distribution.

Let $P(X)$ be the observed distribution. Suppose we have n hypothesized distributions $\{H_j(\alpha_j^l, X) : j = 1, 2, \dots, n\}$, where $\alpha_j^l : l = 1, \dots, m_j$ are the parameters of distribution $H_j(X)$. We compute the ζ distance between H_j and P ,

$$\zeta(P, H_j) = -\log_2 \left[\frac{1 + \sum_k \sqrt{h_j(\alpha_j^l, k)p(k)}}{2} \right], \quad (51)$$

and minimize w.r.t the parameters:

$$\frac{\partial \zeta(P, H_j)}{\partial \alpha_j^l} = 0, \quad l = 1, \dots, m_j. \quad (52)$$

and get the estimated function $\tilde{H}_j(\tilde{\alpha}_j^l, k)$. We need to ensure that this differential equation gives us the minima rather than the maxima of ζ distance. We can then choose the distributions with the minimal ζ distance as the best fit. Instead, if the model parameters are estimated through other means such as maximum likelihood or Bayes estimators, we need additional criterion such as Akaike information criterion (AIC) to test the goodness of fit of models. Here we propose that choosing model with minimal distance between observed and model distribution can serve as an additional criterion. AIC provides an asymptotically unbiased estimator of KLD separation between data and model [34]. It works well when the sample size is large and the number of parameters is comparatively small. KLD is just one measure to assess the disparity between true and candidate model. To the best of our knowledge, we do not find any studies using other measures of disparity such as Hellinger, Bhattacharyya or Jensen-Shannon divergence in AIC type criteria. Here we propose to use these measures along with BBD to assess the disparity between data and model. We believe that these measures might shed additional light into parameter estimation and address limitations of KLD based AIC. We have not tested our proposed method on real data. We leave to future studies to test the robustness of such an approach to model selection and parameter estimation.

VI. CONCLUSION

In this work we have introduced a new bounded divergence measure based on Bhattacharyya distance. It belongs to the class of f-divergences and shares all its characteristics. Although many bounded divergence measures have been studied and used in various applications, no single 'metric' is useful in all types of problems studied. Ours is based on the Bhattacharyya coefficient which is useful in computing tight bounds on Bayes error probabilities. The Bounded Bhattacharyya distance shares many common properties with

Hellinger, Bhattacharyya and Jensen-Shannon divergence measures and we have provided several inequalities relating them. We have also proposed a new method for parameter estimation for probability distributions based on divergence measures. We are investigating further, properties of the BBD measure and plan to apply it to network models and population dynamics.

VII. ACKNOWLEDGEMENTS

One of us (S.J) thanks Rahul Kulkarni for insightful discussions, and acknowledge the financial support in part by grants DMR-0705152 and DMR-1005417 from the US National Science Foundation. M.S. would like to thank the Penn State Electrical Engineering Department for support.

VIII. APPENDIX

A. ζ measures of some common distributions

• Binomial :

$$P(k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad Q(k) = \binom{n}{k} q^k (1-q)^{n-k}.$$

$$\zeta_{bin}(P, Q) = -\log_2 \left(\frac{1 + [\sqrt{pq} + \sqrt{(1-p)(1-q)}]^n}{2} \right). \quad (53)$$

• Poisson :

$$P(k) = \frac{\lambda_p^k e^{-\lambda_p}}{k!}, \quad Q(k) = \frac{\lambda_q^k e^{-\lambda_q}}{k!}.$$

$$\zeta_{poisson}(P, Q) = -\log_2 \left(\frac{1 + e^{-(\sqrt{\lambda_p} - \sqrt{\lambda_q})^2/2}}{2} \right). \quad (54)$$

• Gaussian :

$$P(x) = \frac{1}{\sqrt{2\pi}\sigma_p} \exp \left(-\frac{(x-x_p)^2}{2\sigma_p^2} \right),$$

$$Q(x) = \frac{1}{\sqrt{2\pi}\sigma_q} \exp \left(-\frac{(x-x_q)^2}{2\sigma_q^2} \right).$$

$$\zeta_{Gauss}(P, Q) = 1 - \log_2 \left[1 + \frac{2\sigma_p\sigma_q}{\sigma_p^2 + \sigma_q^2} \exp \left(-\frac{(x_p - x_q)^2}{4(\sigma_p^2 + \sigma_q^2)} \right) \right]. \quad (55)$$

• Exponential : $P(x) = \lambda_p e^{-\lambda_p x}$, $Q(x) = \lambda_q e^{-\lambda_q x}$.

$$\zeta_{exp}(P, Q) = -\log_2 \left[\frac{(\sqrt{\lambda_p} + \sqrt{\lambda_q})^2}{2(\lambda_p + \lambda_q)} \right]. \quad (56)$$

• Pareto : Assuming the same cut off x_m ,

$$P(x) = \begin{cases} \alpha_p \frac{x_m^{\alpha_p}}{x^{\alpha_p+1}} & \text{for } x \geq x_m \\ 0 & \text{if } x < x_m, \end{cases} \quad (57)$$

$$Q(x) = \begin{cases} \alpha_q \frac{x_m^{\alpha_q}}{x^{\alpha_q+1}} & \text{for } x \geq x_m \\ 0 & \text{if } x < x_m. \end{cases} \quad (58)$$

$$\zeta_{pareto}(P, Q) = -\log_2 \left[\frac{(\sqrt{\alpha_p} + \sqrt{\alpha_q})^2}{2(\alpha_p + \alpha_q)} \right]. \quad (59)$$

REFERENCES

- [1] S. Kullback and R. A. Leibler, "On information and sufficiency," *The Annals of Mathematical Statistics*, vol. 22, no. 1, pp. 79–86, 1951.
- [2] S. M. Ali and S. D. Silvey, "A general class of coefficients of divergence of one distribution from another," *Journal of the Royal Statistical Society. Series B (Methodological)*, vol. 28, no. 1, pp. 131–142, 1966.
- [3] J. Kapur, "A comparative assessment of various measures of directed divergence," *Advances in Management Studies*, vol. 3, no. 1, pp. 1–16, 1984.
- [4] S. Kullback, "Information theory and statistics," *New York: Dover*, 1968, 2nd ed., vol. 1, 1968.
- [5] U. Kumar, V. Kumar, and J. N. Kapur, "Some normalized measures of directed divergence," *International Journal of General Systems*, vol. 13, no. 1, pp. 5–16, 1986.
- [6] M. Basseville, "Distance measures for signal processing and pattern recognition," *Signal processing*, vol. 18, pp. 349–369, 1989.
- [7] M. Ben-Bassat, "f-entropies, probability of error, and feature selection," *Information and Control*, vol. 39, no. 3, pp. 227–242, 1978.
- [8] E. Choi and C. Lee, "Feature extraction based on the Bhattacharyya distance," *Pattern Recognition*, vol. 36, no. 8, pp. 1703–1709, Aug 2003.
- [9] T. Kailath, "The Divergence and Bhattacharyya Distance Measures in Signal Selection," *IEEE Transactions on Communications*, vol. 15, no. 1, pp. 52–60, Feb. 1967.
- [10] T. Kadota and L. Shepp, "On the best finite set of linear observables for discriminating two gaussian signals," *IEEE Transactions on Information Theory*, vol. 13, no. 2, pp. 278–284, 1967.
- [11] K. Tumer and J. Ghosh, "Estimating the Bayes error rate through classifier combining," *Proceedings of 13th International Conference on Pattern Recognition*, pp. 695–699, 1996.
- [12] M. Nielsen and I. Chuang, "Quantum computation and information," *Cambridge University Press, Cambridge, UK*, vol. 3, no. 8, p. 9, 2000.
- [13] P. W. Lamberti, A. P. Majtey, A. Borras, M. Casas, and A. Plastino, "Metric character of the quantum Jensen-Shannon divergence," *Physical Review A*, vol. 77, p. 052311, 2008.
- [14] A. Gibbs and F. Su, "On choosing and bounding probability metrics," *International Statistical Review*, vol. 70, no. 3, pp. 419–435, 2002.
- [15] A. Bhattacharyya, "On a measure of divergence between two multinomial populations," *Sankhyā: The Indian Journal of Statistics (1933-1960)*, vol. 7, no. 4, pp. 401–406, 1946.
- [16] F. Nielsen and S. Boltz, "The burbea-rao and bhattacharyya centroids," *Information Theory, IEEE Transactions on*, vol. 57, no. 8, pp. 5455–5466, 2011.
- [17] H. Chernoff, "A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations," *The Annals of Mathematical Statistics*, vol. 23, no. 4, pp. 493–507, 1952.
- [18] F. Nielsen, "Chernoff information of exponential families," *Arxiv preprint arXiv:1102.2684*, 2011.
- [19] A. DasGupta, *Probability for Statistics and Machine Learning*, ser. Springer Texts in Statistics. Springer New York, 2011.
- [20] J. Burbea and C. R. Rao, "On the convexity of some divergence measures based on entropy functions," *IEEE Transactions on Information Theory*, vol. 28, no. 3, pp. 489 – 495, may 1982.
- [21] C. R. Rao, "Diversity and dissimilarity coefficients: A unified approach," *Theoretical Population Biology*, vol. 21, no. 1, pp. 24 – 43, 1982.
- [22] J. Lin, "Divergence measures based on the shannon entropy," *IEEE Transactions on Information Theory*, vol. 37, no. 1, pp. 145 –151, jan 1991.
- [23] A. Bhattacharyya, "On a measure of divergence between two statistical populations defined by their probability distributions," *Bull. Calcutta Math. Soc.*, vol. 35, no. 99-109, p. 4, 1943.
- [24] E. Hellinger, "Neue begründung der theorie quadratischer formen von unendlichvielen veränderlichen," *Journal für die reine und angewandte Mathematik (Crelle's Journal)*, vol. 1909, no. 136, pp. 210–271, 1909.
- [25] S. Kakutani, "On equivalence of infinite product measures," *The Annals of Mathematics*, vol. 49, no. 1, pp. 214–224, 1948.
- [26] M. E. Hellman and J. Raviv, "Probability of Error, Equivocation, and the Chernoff Bound," *IEEE Transactions on Information Theory*, vol. 16, no. 4, pp. 368–372, 1970.

- [27] R. Bradt and S. Karlin, "On the design and comparison of certain dichotomous experiments," *The Annals of mathematical statistics*, pp. 390–409, 1956.
- [28] D. Blackwell, "Comparison of experiments," in *Second Berkeley Symposium on Mathematical Statistics and Probability*, vol. 1, 1951, pp. 93–102.
- [29] I. Csiszar, "Information-type distance measures and indirect observations," *Stud. Sci. Math. Hungar.*, vol. 2, pp. 299–318, 1967.
- [30] —, "I-divergence geometry of probability distributions and minimization problems," *The Annals of Probability*, vol. 3, no. 1, pp. 146–158, 1975.
- [31] R. Sibson, "Information radius," *Probability Theory and Related Fields*, vol. 14, no. 2, pp. 149–160, 1969.
- [32] C. R. Rao, "Information and the accuracy attainable in the estimation of statistical parameters," *Bull. Calcutta Math. Soc.*, vol. 37, pp. 81–91, 1945.
- [33] —, "Differential metrics in probability spaces," *Differential geometry in statistical inference*, vol. 10, pp. 217–240, 1987.
- [34] K. Burnham and D. Anderson, *Model selection and multimodel inference: a practical information-theoretic approach*. Springer Verlag, 2002.